

# Nested Cantor sets

Pierre Berger <sup>\*</sup> and Carlos Gustavo Moreira <sup>†</sup>

March 6, 2013

## Abstract

We give sufficient conditions for two Cantor sets of the line to be nested for a positive set of translation parameters. This problem is a toy model of the parameter selection for non-uniformly hyperbolic attractors of the plane. In natural such models, we show that this condition is optimal.

## Introduction

One dimensional Cantor sets of positive measure appear frequently in dynamical systems. For instance, if a quadratic map  $f(x) = x^2 + a$  has an absolutely continuous invariant measure (acim) with Lyapunov exponent  $\lambda > 0$ , then for  $C > 0$  small enough,  $\lambda' \in (0, \lambda)$ , the set  $\tilde{K}$  of points  $x$  such that:

$$(\star) \quad \|D_x f^n\| \geq C\lambda'^n, \quad \forall n \geq 0.$$

is a Cantor set of positive Lebesgue measure. Collet-Eckmann property is that the critical value belongs to such a  $\tilde{K}$  (for a certain  $C, \lambda'$ ). Actually the set of Collet-Eckmann parameters  $a$  is of positive measure [BC89, Yoc97]. A rough idea of the proof is the following: the critical value  $a$  “moves faster” than the set  $\tilde{K}$ .

In higher dimension, for instance in the study of the Hénon map [Ber11], a non-uniformly hyperbolic attractor appears when a cantor set  $K$  is included in a similar Cantor set similar to  $\tilde{K}$ . A main issue is to find such non-uniformly attractors with a dimension  $d$  not too small, for a positive set of parameters of generic family of maps. As a toy model for the parameter selection, we can assume that  $K$  and  $\tilde{K}$  do not depend on the parameter. This leads to the following question:

**Question 0.1** ([Ber11]). *Under which hypothesis on a Cantor set  $K$  and a Lebesgue positive Cantor set  $\tilde{K}$ , the set of parameters  $t \in \mathbb{R}$  such that  $K+t$  is included in  $\tilde{K}$  has a positive Lebesgue measure?*

---

<sup>\*</sup>berger@math.univ-paris13.fr, LAGA Université Paris 13, partially financed by the Balzan prize of J. Palis

<sup>†</sup>gugu@impa.br, Instituto de Matemática Pura e Aplicada; partially financed by the Balzan prize of J. Palis and by CNPq

**Remark 0.2.** If  $K$  is of positive measure and  $t + K$  is included  $\tilde{K}$  for a Lebesgue positive set  $T$  of parameters  $t$ , then  $\tilde{K}$  must contains an interval, which is impossible since it is a Cantor set.

In other words, if  $K$  and  $T$  have positive measure,  $K + T \subset \tilde{K}$  has non-empty interior. Indeed,  $K + T = K - (-T) = \{s | K \cap (s - T) \neq \emptyset\}$ . Since  $K$  and  $-T$  have positive measure, there is  $\epsilon > 0$  and intervals  $[a, a + \epsilon]$  and  $[b, b + \epsilon]$  such that  $\text{Leb}([a, a + \epsilon] \cap K) > 3\epsilon/4$  and  $\text{Leb}([b, b + \epsilon] \cap (-T)) > 3\epsilon/4$ . So, if  $s \in (b - a - \epsilon/2, b - a + \epsilon/2)$  then  $K \cap (s - T) \neq \emptyset$ .

We will gives a sufficient condition for a positive answer to this question, based on the following condition on  $\tilde{K}$ .

**Definition 0.3.** For  $0 < p < 1$ , a compact set  $\tilde{K} = [a, a + \text{diam } \tilde{K}] \setminus \sqcup_n (a_n, b_n)$ , where the intervals  $(a_n, b_n)$  are the connected components of  $[a, a + \text{diam } \tilde{K}] \setminus \tilde{K}$  satisfies condition  $(\mathcal{C}_p)$  if:

$$(\mathcal{C}_p) \quad \sum_n (b_n - a_n)^p < \infty.$$

We define

$$P(\tilde{K}) := \inf\{p > 0 : \tilde{K} \text{ satyisfies condition } (\mathcal{C}_p)\}$$

Let us formulate a few remarks on condition  $(\mathcal{C}_p)$ .

**Remark 0.4.** If the complement of  $\tilde{K}$  is expressed as an union of intervals which are not necessarily connected components of  $[a, a + \text{diam } \tilde{K}] \setminus \tilde{K}$ :

$$\tilde{K} = [a, a + \text{diam } \tilde{K}] \setminus \cup_n (a_n, b_n),$$

such that  $\sum_n (b_n - a_n)^p < \infty$ , then  $\tilde{K}$  satisfies condition  $(\mathcal{C}_p)$ . Indeed, for every positive numbers  $(l_i)_i$ , it holds  $\sum_i l_i^p \geq (\sum_i l_i)^p$ , hence the sum  $\sum_n (b_n - a_n)^p$  is smaller if we group the intervals in the same components.

The inequality  $\sum_i l_i^p \geq (\sum_i l_i)^p$  also implies the following

**Remark 0.5.** If  $\tilde{K}_1$  and  $\tilde{K}_2$  are two subsets of  $[0, 1]$  satisfying  $(\mathcal{C}_p)$ , then  $\tilde{K}_1 \cap \tilde{K}_2$  satisfies  $(\mathcal{C}_p)$ .

A consequence of the main result is the following:

**Proposition 0.6.** *Let  $K$  be a regular Cantor set of Hausdorff dimension  $d$  of  $\mathbb{R}$  and let  $\tilde{K}$  be a Cantor set of positive measure satisfying condition  $(\mathcal{C}_p)$ . If  $1 - d > p$  then for  $\lambda > 0$  sufficiently small, the following set:*

$$\{t \in \mathbb{R} : t + \lambda K \subset \tilde{K}\},$$

*has positive Lebesgue measure.*

Main Theorem 2.1 gives a sufficient condition to answer to question 0.1. It implies the above Proposition.

We define in section 1 the different notions used by Proposition 0.6 and Theorem 2.1. In section 2, we state the main theorem and prove Proposition 0.6. Then we apply this Proposiotion to many examples which are toy models for the parameter selection of non-uniformly hyperbolic attractor. In section 3, Theorem 2.1 is proven. In section 4, we discuss on the optimality of the result, thanks to example and counter example, some of which are toy models for the parameter selection of non-uniformly hyperbolic attractor. In section 5, we will explain the above toy model.

# 1 Notations and definitions

## 1.1 Operations on sets

Given two subsets  $E, F \subset \mathbb{R}$ , the arithmetic sums  $E + F$  and  $E - F$  stand for the following subsets of  $\mathbb{R}$ :

$$E + F = \{e + f : e \in E, f \in F\}, \quad E - F = \{e - f : e \in E, f \in F\}$$

We denote also by  $-E$  the set  $\{0\} - E$ .

The operations  $+$  and  $-$  are associative. Thus, for any three subsets  $E, F, G \subset \mathbb{R}$ , we can denote by  $E + F + G$  the set:

$$E + F + G = E + (F + G) = (E + F) + G.$$

For  $U \in \mathbb{R}$ , let  $U \cdot E := \{U \cdot e : e \in E\}$ . However  $2 \cdot E \subsetneq E + E$  if  $E$  has more than one element.

## 1.2 Regular Cantor sets

**Definition 1.1** (Regular Cantor set). We recall that  $K$  is a  $C^k$ -regular Cantor set,  $k \geq 1$ , if:

- i) there are disjoint compact intervals  $I_1, I_2, \dots, I_r$  such that  $K \subset I_1 \cup \dots \cup I_r$  and the boundary of each  $I_j$  is contained in  $K$ ;
- ii) there is a  $C^k$  expanding map  $\psi$  defined in a neighbourhood of  $I_1 \cup I_2 \cup \dots \cup I_r$  such that  $\psi(I_j)$  is the convex hull of a finite union of some intervals  $I_s$  satisfying:
  - ii.1) for each  $j$ ,  $1 \leq j \leq r$  and  $n$  sufficiently big,  $\psi^n(K \cap I_j) = K$ ;
  - ii.2)  $K = \bigcap_{n \in \mathbb{N}} \psi^{-n}(I_1 \cup I_2 \cup \dots \cup I_r)$ .

There are two basic concepts of dimension for Cantor sets:

**Definition 1.2** (Box dimension and Box fuzzy measure). The *box dimension*  $d$  of a compact set  $K$  is the infimum of  $s \geq 0$  such that the following is zero:

$$\limsup_{\epsilon \rightarrow 0} \inf \{N\epsilon^s : (U_i)_{i=1}^N \text{ is a finite covering of } K \text{ by open intervals of diameter } \epsilon\}$$

We denote by  $C_K$  the following supremum, with  $d$  the box dimension:

$$C_K := \sup_{0 < \epsilon \leq \text{diam } K} \inf \{N\epsilon^d : (U_i)_{i=1}^N \text{ is a finite covering of } K \text{ by open intervals of diameter } \epsilon\}$$

The map  $K \mapsto C_K$  is a fuzzy measure, but not a measure. We call it the *d-box fuzzy measure*.

**Remark 1.3.** For every compact set  $K$  of box dimension  $d$  and  $\lambda > 0$ , it holds:

$$\text{diam } (\lambda K) = \lambda \text{diam } K \quad \text{and} \quad C_{\lambda K} = \lambda^d C_K.$$

The box dimension is at least equal to the Hausdorff dimension:

**Definition 1.4** (Hausdorff dimension). The *Hausdorff dimension*  $d$  of a compact set  $K$  is the infimum of  $s$  such that the following is zero:

$$\limsup_{\epsilon \rightarrow 0} \inf \left\{ \sum_i \text{diam}(U_i)^s : (U_i)_i \text{ is a covering of } K \text{ by intervals of diameter } \leq \epsilon \right\}$$

The value of the above limit at  $s = d$  is denoted by  $m_d(K)$ . The map  $K \mapsto m_d(K)$  is a measure, called the Hausdorff  $d$ -dimensional measure.

In the case of regular Cantor set, these two definitions coincide from the following.

**Theorem 1.5** (Palis-Takens Prop. 3 p. 72 [PT93]). *If  $K$  is a regular Cantor set of class  $C^{1+\alpha}$ , the Hausdorff dimension  $d$  of  $K$  is equal to the Box dimension of  $K$ . Moreover, the box fuzzy measure  $C_K$  is finite.*

## 2 Main result and its applications

Here is the main result of this article. Its proof will be done in §3.

**Theorem 2.1.** *Let  $\tilde{K} = [0, \text{diam } \tilde{K}] \setminus \sqcup_n (a_n, b_n)$  be a Cantor set satisfying condition  $(\mathcal{C}_p)$ . Put  $l_n := b_n - a_n$ . Let  $K$  be a regular Cantor set of dimension  $d < 1 - p$  and box fuzzy measure  $C_K$  satisfying:*

$$\sum_{\{n: l_n > \text{diam } K\}} (\text{diam } K + l_n) + 2C_K \sum_{\{n: l_n \leq \text{diam } K\}} (l_n)^{1-d} < \text{diam}(\tilde{K}) - \text{diam}(K),$$

*Then there exists a set of Lebesgue positive measure set of parameters  $t \in \mathbb{R}$  such that  $t + K$  is included in  $\tilde{K}$ .*

*Proof of Proposition 0.6.* Without loss of generality we can suppose that the diameter of  $K$  is 1, and hence for  $\lambda > 0$ :

$$\text{diam}(\lambda K) = \lambda$$

We recall that for every regular Cantor set  $K$  of dimension  $d$  and  $\lambda > 0$ , it holds:

$$C_{\lambda K} = \lambda^d C_K$$

By Theorem 2.1, Proposition holds if for  $\lambda$  small:

$$(1) \quad \sum_{\{n: l_n > \lambda\}} (\lambda + l_n) + 2\lambda^d C_K \sum_{\{n: l_n \leq \lambda\}} (l_n)^{1-d} < \text{diam}(\tilde{K}) - \lambda,$$

where  $(l_n)_n$  are the diameter of the holes of  $\tilde{K}$ . As condition  $(\mathcal{C}_p)$ ,  $p < 1 - d$  is satisfied by  $\tilde{K}$ , inequality (1) holds if there exists  $\eta > 0$  such that for  $\lambda$  small:

$$(2) \quad \sum_{\{n: l_n > \lambda\}} (\lambda + l_n) < \text{diam}(\tilde{K}) - \eta.$$

We remark that for every  $N \geq 0$ :

$$\sum_{\{n: l_n > \lambda\}} (\lambda + l_n) = \sum_{\{n \leq N: l_n > \lambda\}} (\lambda + l_n) + \sum_{\{n > N: l_n > \lambda\}} (\lambda + l_n) \leq \sum_{\{n \leq N: l_n > \lambda\}} (\lambda + l_n) + \sum_{n > N} 2l_n.$$

For  $N$  large the second sum is small, and then by taking  $\lambda$  small (in function of  $N$ ), the first sum is close to  $\sum_n (l_n) = \text{Leb}([0, \text{diam } \tilde{K}] \setminus \tilde{K})$ . Consequently inequality (2) holds for  $\lambda$  small if

$$\text{Leb}([0, \text{diam } \tilde{K}] \setminus \tilde{K}) < \text{diam } (\tilde{K})$$

which is indeed the case since  $\tilde{K}$  has positive Lebesgue measure.  $\square$

Let us apply Theorem 2.1 and Proposition 0.6 to various examples issued from non uniformly hyperbolic dynamics.

**Example 2.2.** The simplest example of non uniformly hyperbolic map is the Chebichev map  $x \mapsto x^2 - 2$ . It is semi conjugated to the doubling angle maps  $\theta$  via a map which have a quadratic singularity. Hence a first toy model of the set of points with positive Lyapunov exponent is the following Cantor set  $\tilde{K}$  with  $s < 1/2$ , where we cut the circle at a point to make the segment  $[-1, 1]$ .

Let  $\tilde{K}(s)$  be the Cantor set  $\cap_{n \geq N} \tilde{K}_n$  where  $\tilde{K}_n$  is an union of  $2^n$  intervals constructed by induction as follows:

- $\tilde{K}_n := [-1, 1]$ , for  $n \leq [\log_2(1 - 2^{-s})/s] =: n_0$ .
- For  $N > n_0$ ,  $\tilde{K}_n$  is obtained by taking away from each component of  $\tilde{K}_{n-1}$  an open interval of length  $2^{-n/s}$ , for instance in the middle of the component.

The complement of  $[-1, 1] \setminus \tilde{K}(s)$  has  $2^{n-n_0}$  components of length  $2^{-n/s}$ . Hence this Cantor set has positive Lebesgue measure and Condition  $(\mathcal{C}_p)$  is satisfied for  $p > s$  (and not satisfied for  $p \leq s$ ), that is,  $P(K(s)) = s$ .

Hence by Proposition 0.6 the Cantor set  $\tilde{K}(s)$  contains a positive set of translations of a dyadic Cantor set  $K_d$  of dimension  $d < 1 - s$ .

**Example 2.3.** Let us modelize better the Chebichev map  $x \mapsto x^2 - 2$  by looking its first return map into  $[-1, 1]$ . Such a first return map is smoothly conjugated to the following map:

$$f: x \in [-1, 1] \setminus \{0\} \mapsto \begin{cases} 2^{n+2}(x - 2^{-n-1}) - 1, & \text{if } x \in (2^{-n-1}, 2^{-n}] \\ 2^{n+2}(-x - 2^{-n-1}) - 1, & \text{if } -x \in (2^{-n-1}, 2^{-n}] \end{cases}$$

Put  $I_{n+2} := (2^{-n-1}, 2^{-n}]$  and  $I_{-n-2} := [-2^{-n}, -2^{-n-1})$ . Both intervals  $\text{int } I_{\pm(n+2)}$  are sent diffeomorphically onto  $(-1, 1)$  by  $f$ . They are expanded by a factor  $2^{n+2}$ .

The conjugacy with the Chebichev map preserves 0, hence a point in  $I_n$  will be contracted by a factor of order  $2^{-n}$ .

Let  $\mathcal{D} := \cap_{k \geq 0} f^{-k}([-1, 1])$  which is equal to  $[-1, 1]$  but a countable set.

Every  $x \in \mathcal{D}$  has its image by  $f^k$  into a certain  $I_{x_k}$ , with  $x_k \in \mathbb{Z} \setminus \{-1, 0, 1\}$ . Using the bijection map  $x \in \mathcal{D} \mapsto (x_k)_{k \geq 1}$ , we define the following family of Cantor sets:

$$\tilde{K}_2(s, N) := cl(\{x \in \mathcal{D} : |x_i| \leq \max(N, s \sum_{j < i} |x_j|), \quad \forall i\}).$$

We remark that the set of points with positive Lyapunov exponent for the Chebichev map is diffeomorphic to:

$$\bigcup_{s < 1, N \geq 0} \tilde{K}_2(s, N)$$

Let us compute what is the optimal condition  $(\mathcal{C}_p)$  satisfied by  $\tilde{K}_2(s, N)$ . Since  $|f'|_{I_k} \equiv 2^{-|k|}$ ,  $\cap_{j=0}^{i-1} f^{-j}(I_{x_j})$  is an interval  $\tilde{I}$  of size  $2^{1-S}$ , where  $S = \sum_{0 \leq j < i} |x_j|$ , and, for  $i$  large, the condition  $|x_i| > \max(N, s \sum_{j < i} |x_j|)$  is equivalent to  $|x_i| > s \sum_{j < i} |x_j| = sS$ , and so corresponds to a gap in the middle of  $\tilde{I}$  and proportion  $2^{-s \sum_{j < i} |x_j|} = 2^{-sS}$ , so with size  $2(|\tilde{I}|/2)^{-(1+s)}$ . These are (except for the small values of  $i$ ) the connected components of the complement of  $\tilde{K}_2(s, N)$ . So, analogously as the previous example,  $P(\tilde{K}_2(s, N)) = \frac{1}{1+s}$ .

Actually, in the parameter selection of non uniformly hyperbolic attractors, the intervals  $I_n$ , for  $n$  large, are constructed inductively from the intervals in which the critical points went through. In such operation the hyperbolicity is lost, and so the critical point should not pass, in mean, by too much intervals  $I_n$ , for  $n$  large. This gives rise to the notion of strong regularity [Yoc97, Ber11].

**Example 2.4.** Using the above coding of  $\mathcal{D}$ , for  $M > 0$ , let:

$$\tilde{K}_M := \{x \in [-1, 1] \text{ s.t. } \forall k \geq 0, \quad \sum_{j \leq k+1: |x_j| > M} |x_j| \leq 2^{-\sqrt{M}} \sum_{j \leq k} |x_j|\}$$

This set corresponds, in the dictionary puzzle-parapuzzle, to the set of strongly regular quadratic maps [Yoc97]. These quadratic maps satisfy the Collet-Eckman condition. Let us suppose that  $\tilde{K}_M$  satisfies  $(\mathcal{C}_p)$ . Let us bound from below the value of  $p$ .

It is well known that, for any positive  $M$ ,  $\tilde{K}_M$  has positive Lebesgue measure, and so there is  $c > 0$  such that, for infinitely many values of  $n$ , the number of intervals of the type  $\{x : |x_1| = a_1, |x_2| = a_2, \dots, |x_i| = a_i\}$  which have size  $2^{-n}$  and intersect  $\tilde{K}_M$  is at least  $c \cdot 2^n$ . For these intervals, the intervals

$$\{x : |x_1| = a_1, |x_2| = a_2, \dots, |x_i| = a_i, \quad |x_{i+1}| > 2^{-\sqrt{M}} n\}$$

are components of the complement of  $(\tilde{K}_M)$ , and their length is equivalent to  $2^{-n} 2^{-n \cdot 2^{-\sqrt{M}} + 1} = 2^{1-(1+2^{-\sqrt{M}})n}$ .

Consequently, if Condition  $(\mathcal{C}_p)$  is satisfied, the following sum converges:

$$\sum_n 2^{n-p(1+n \cdot 2^{-\sqrt{M}})+p}$$

This happens iff  $1 < p(1 + 2^{-\sqrt{M}})$ , that is:

$$p > \frac{1}{1 + 2^{-\sqrt{M}}}.$$

So,  $P(\tilde{K}_M) \geq \frac{1}{1+2^{-\sqrt{M}}}$ .

In order to give an upper bound for  $P(\tilde{K}_M)$ , we will assume  $M$  large enough. Let  $\delta = 2^{-\sqrt{M}}$ . The set  $[-1, 1] \setminus \tilde{K}_M$  is the union of intervals of the type

$$\{x : |x_1| = a_1, |x_2| = a_2, \dots, |x_{n-1}| = a_{n-1}, |x_n| > a_n\}$$

where  $a_n \geq M$ ,  $\sum_{j \leq n: |a_j| > M} |a_j| > \delta \sum_{j \leq n-1} |a_j|$ , and so, for  $N = \sum_{j \leq n} |a_j| \geq 2n$ , the size of this interval is  $O(2^{-N})$ , and we have  $\sum_{j \leq n: |a_j| > M} |a_j| > \delta N$ . Let  $R = \sum_{j \leq n: |a_j| > M} |a_j| > \delta N$ . If  $t = |\{j \leq n : |a_j| > M\}|$ , we have  $t \leq R/M$ , and the number of possible choices of  $\{j \leq n : |a_j| > M\}$  is  $O(\binom{n}{t}) = O(\binom{n}{R/M}) = O((enM/R)^{R/M}) = O((eNM/2R)^{R/M})$  and the number of possible choices of  $(a_j, j \leq n : |a_j| > M)$  is bounded by  $2^t$  times the number of solutions of  $x_1 + x_2 + \dots + x_t = R$ , so it is  $O(2^t \binom{R}{t}) = O(2^t \binom{R}{R/M}) = O((2eM)^{R/M})$ . The number of choices of  $(a_j, j \leq n : |a_j| \leq M)$  is bounded by  $2^{N-R}$ . So, the total number of such intervals for given  $N, t, R$  is  $O((e^2 M^2 N/R)^{R/M} \cdot 2^{N-R}) = O(2^{N-R} (e^2 M^2 / \delta)^{R/M}) = O(2^{N-R} (e^2 M^2 2^{\sqrt{M}})^{R/M}) = O(2^N (e^2 M^2 2^{\sqrt{M}-M})^{R/M}) = O(2^N (e^2 M^2 2^{\sqrt{M}-M})^{2^{-\sqrt{M}} N/M})$ . Given  $N$  there are at most  $N^2$  choices for  $(t, R)$ , and so the total number of possibilities is  $O(2^N \cdot N^2 (e^2 M^2 2^{\sqrt{M}-M})^{2^{-\sqrt{M}} N/M}) = O(2^N (2^{2\sqrt{M}-M})^{2^{-\sqrt{M}} N/M}) = O(2^{N(1-(\frac{2}{\sqrt{M}}-1)2^{-\sqrt{M}})})$ . Since the size of the corresponding interval is  $O(2^{-N})$ , this implies that  $P(\tilde{K}_M) \leq 1 - (\frac{2}{\sqrt{M}} - 1)2^{-\sqrt{M}}$ .

### 3 Proof of Theorem 2.1

We recall that  $\tilde{K}$  is included in  $[0, \text{diam } \tilde{K}]$  with  $[0, \text{diam } \tilde{K}] \setminus \tilde{K} = \sqcup_n (a_n, b_n)$ .

We can suppose that the regular Cantor set  $K$  of dimension  $d$  is included in  $[0, \text{diam } K]$ . Observe that if  $t$  is in  $[0, \text{diam } \tilde{K} - \text{diam } K]$ , then  $t + K$  is included in  $[0, \text{diam } \tilde{K}]$ .

Let  $\mathbb{X}$  be the set of parameters  $t$  such that  $t + K \subset \tilde{K}$ .

Let us consider  $t \in [0, \text{diam } \tilde{K} - \text{diam } K] \setminus \mathbb{X}$ , in other words  $K + t$  is not included in  $\tilde{K}$ .

This means that  $K + t$  intersects the interval  $(a_n, b_n)$  for a certain  $n$ . That is to say:  $t \in (a_n, b_n) - K$ . This implies:

**Fact 3.1.**  $[0, \text{diam } \tilde{K} - \text{diam } K] \setminus \mathbb{X} = \bigcup_n ((a_n, b_n) - K)$ .

A direct consequence is the following nice inequality:

**Corollary 3.2.**  $\text{Leb}([0, \text{diam } \tilde{K} - \text{diam } K] \setminus \mathbb{X}) \leq \sum_n \text{Leb}((a_n, b_n) - K)$ .

The measure of  $(a_n, b_n) - K$  is estimated by the following Proposition shown below.

**Proposition 3.3.** *For every interval  $I$  of  $\mathbb{R}$ , we have:*

$$\begin{cases} \text{Leb}(I - K) \leq 2C_K (\text{Leb } I)^{1-d} & \text{if } \text{diam } I \leq \text{diam } K \\ \text{Leb}(I - K) \leq \text{Leb } I + \text{diam } K & \text{if } \text{diam } I \geq \text{diam } K \end{cases}$$

with  $C_K$  the box fuzzy measure of dimension  $d$ .

From Proposition 3.2, with  $l_n = b_n - a_n$ :

$$\begin{aligned} \text{Leb}([0, \text{diam } \tilde{K} - \text{diam } K] \setminus \mathbb{X}) &\leq \sum_n \text{Leb}((a_n, b_n) - K) \\ &\leq \sum_{\{n: l_n > \text{diam } K\}} (\text{diam } K + l_n) + \sum_{\{n: l_n \leq \text{diam } K\}} 2C_K l_n^{1-d}, \end{aligned}$$

which implies Theorem 2.1.

*Proof of Proposition 3.3.* To see the case  $\text{diam } I \geq \text{diam } K$  consider that  $K \supset [0, \text{diam } K]$  and that  $I = [-\text{diam } I, 0]$ , which can be considered using a translation. Then  $K - I = [0, \text{diam } K + \text{diam } I]$ .

Let us study the case  $\text{diam } (I) \leq \text{diam } (K)$ . We can assume  $I = [-\epsilon/2, \epsilon/2]$  with  $\epsilon = \text{Leb } I$ .

First observe that

$$(3) \quad [-\epsilon/2, \epsilon/2] - K = -(K - [-\epsilon/2, \epsilon/2]) = -(K + [-\epsilon/2, \epsilon/2]).$$

Note that  $K + [-\epsilon/2, \epsilon/2]$  is the closure of the  $\frac{\epsilon}{2}$ -neighborhood of  $K$ .

By definition of  $C_K$ , the Cantor set  $K$  is included in the union of  $C_K \epsilon^{-d}$  intervals of length  $\epsilon$ . The  $\epsilon/2$ -neighborhood of this union is contained in the union  $U$  of  $C_K \epsilon^{-d}$  intervals of length  $2\epsilon$ , the Lebesgue measure of which is

$$\text{Leb}(U) \leq 2C_K \epsilon^{1-d} = 2C_K (\text{Leb } I)^{1-d}.$$

As  $U$  contains  $K + [-\epsilon/2, \epsilon/2]$ , this proves the Proposition.  $\square$

## 4 Discussion on the optimality of Condition $(\mathcal{C}_p)$

### 4.1 Example for which Proposition 0.6 is trivially non optimal

Condition  $(\mathcal{C}_p)$  is not optimal for Theorem 2.1. This follows from the following observations.

Consider two Cantor sets  $\tilde{K}_1$  and  $\tilde{K}_2$  of positive Lebesgue measure and satisfying respectively  $\mathcal{C}_{p_1}$  and  $\mathcal{C}_{p_2}$ .

The disjoint union  $\tilde{K}$  of two Cantor sets  $\tilde{K}_1$  and  $\tilde{K}_2$  is of positive Lebesgue measure and satisfies, in general,  $\mathcal{C}_p$  only for  $p \geq \max(p_1, p_2)$ , whereas Proposition 0.6 applied to  $\tilde{K}_1$  and  $\tilde{K}_2$  implies that it contains a Lebesgue positive set of translations of a Cantor set with Hausdorff dimension  $1 - \min(p_1, p_2)$ .

In the seek of an optimal result, the following definition makes sens.

**Definition 4.1.** Let  $\tilde{K}$  be a Cantor set of positive measure included in  $[0, 1]$ . Put

$$\hat{P}(\tilde{K}) := \inf\{P(\hat{K}) : \hat{K} \subset \tilde{K} \text{ compact subset of Lebesgue positive measure}\}$$

**Question 4.2.** If  $\tilde{K}$  is a Cantor set of positive measure and containing a positive set of translations of a dyadic Cantor set of dimension  $d$ . Does  $d \leq 1 - P(\tilde{K})$ ?

## 4.2 Negative answer to question 4.2

### 4.2.1 A counter example

For  $p > 0$ , let  $N_p$  be a large integer defined afterward and let  $\tilde{K}_p$  be the Cantor set obtained by taking away of  $[0, 1]$  the following intervals:

$$U_{iq} := \frac{q}{10^i} + (0, 10^{-i/p}), \quad i \geq N_p, \quad q \in \{0, \dots, 10^i - 1\}.$$

We remark that  $\tilde{K}$  has positive measure whenever  $p < 1$  and  $N_p$  is sufficiently large. Moreover  $\tilde{K}_p$  satisfies condition  $\mathcal{C}_{p'}$  for every  $p' > p$ .

The intervals  $(U_{iq})_{iq}$  are nested or disjoint, even if we remove those which are nested,  $\tilde{K}$  does not satisfy Condition  $(\mathcal{C}_{p'})$ , for  $p' \leq p$ .

We remark that  $U_{iq}$  is formed by points  $u$  with the following decimal expression:

$$u = 0, u_1 u_2 \cdots u_i \underbrace{0 \cdots 0}_{(j-i) \times} u_{j+1} \cdots u_n \cdots, \quad (u_l)_l \in \{0, \dots, 9\}$$

such that  $q = \sum_{l=1}^i u_l 10^{i-l}$ ,  $j = [i/p]$ , and  $u_{j+1} \cdots u_n \cdots$  any numbers in  $\{0, \dots, 9\}$  (not all equal to zero).

Let  $K$  be the regular Cantor set of  $[0, 1]$  formed by the points  $k = 0, 0k_1 k_2 \cdots k_j \cdots$  such that  $k_i$  are even number in  $\{0, \dots, 8\}$ . The Hausdorff dimension of such a Cantor set is  $d = \ln(5)/\ln(10)$ . The diameter of  $K$  is less than  $1/10$ .

From fact 3.1, the set  $\mathbb{X}$  of parameters  $t$  such that  $t + K \subset \tilde{K}$  is included in  $[0, \text{diam } \tilde{K} - \text{diam } K]$  and satisfies:

$$[0, \text{diam } \tilde{K} - \text{diam } K] \setminus \mathbb{X} = \bigcup_{k \in K, i \geq N_p} \bigcup_{0 \leq q \leq 10^i} \left\{ \frac{q}{10^i} - k \right\} + (0, 10^{-j}),$$

It is an union of intervals of same length. Let us observe that for every  $p$  the measure of this set is smaller than  $\text{diam } \tilde{K} - \text{diam } K$ , then  $t + K \subset \tilde{K}$  for a Lebesgue positive set of parameters  $t$ .

Let  $K_i$  be the subset of points in  $K$  formed by numbers  $k$  such that  $10^j k$  has fractional part 0. Note that

$$[0, \text{diam } \tilde{K} - \text{diam } K] \setminus \mathbb{X} \subset \bigcup_i \bigcup_{0 \leq q < 10^i, k \in K_j} \{q/10^i - k\} + [-10^{-j}, 10^{-j}].$$

Let us bound from above the cardinality  $C_i$  of  $\{q/10^i - k; 0 \leq q < 10^i, k \in K_j\}$ . Every  $k \in K_j$  is of the form  $k = k_1 + 10^{-i} k_2$  with  $k_1 \in K_i$  and  $k_2 \in K_{j-i}$ . Also

$$\{q/10^i - k; 0 \leq q < 10^i, k_1 \in K_i\} \subset 10^{-i} \cdot \{-10^i + 1, \dots, 0, \dots, 10^i - 1\}$$

which is of cardinality  $2 \cdot 10^i - 1$ . On the other hand,  $K_{j-i}$  is of cardinality  $10^{d(j-i)}$ . Hence:

$$C_i \leq 2 \cdot 10^i \cdot 10^{d(j-i)}$$

Consequently:

$$\text{Leb}([0, \text{diam } \tilde{K} - \text{diam } K] \setminus \mathbb{X}) \leq \sum_{i \geq N_p} C_i 2 \cdot 10^{-j} \leq \sum_{i \geq N_p} 2 \cdot 10^i \cdot 10^{d(j-i)} 2 \cdot 10^{-j} = 4 \sum_{i \geq N_p} 10^{(d-1)([i/p]-i)}.$$

Hence for  $p < 1$ , there exists  $N_p$  such that  $\text{Leb}([0, \text{diam } \tilde{K} - \text{diam } K] \setminus \mathbb{X})$  has measure smaller than  $\text{Leb}([0, \text{diam } \tilde{K} - \text{diam } K]) \leq 0.9$ . Hence  $\mathbb{X}$  has positive measure, and  $K + t$  is contained in  $\tilde{K}_p$  for a Lebesgue positive set of parameters  $t$ .

**Remark 4.3.** This example shows that Theorem 2.1 is not optimal since we can chose  $p$  so that  $p + d > 1$ .

#### 4.2.2 Generalization of the counter example

For the sake of simplicity, we have chosen  $K$  with decimal expression formed only by even numbers, but we could have chosen any other subset  $J$  of  $\{0, \dots, 9\}$ , and have the same result. Observe that the Hausdorff dimension of  $K$  is then  $\ln(\text{card } J)/\ln(10)$ . Moreover, we could have chosen any other basis than the decimal one. This leads to the following observation. For any  $d < 1$ , there exists a regular Cantor set  $K$  of dimension greater than  $d$ , such that for any  $p < 1$ , there exists a Cantor set  $\tilde{K}$  which do not satisfies  $(\mathcal{C}_p)$  but which contains  $K$  for a positive set of translation parameters.

Nevertheless, this counter example to the optimality of Theorem 2.1 does not seem to be stable by perturbation, as shown by the example in §4.3.

#### 4.2.3 Negative answer to question 4.2

Let us come back to the example of §4.2.1.

Let  $\hat{K} \subset \tilde{K}$  be of Lebesgue positive measure. Suppose that  $\hat{K}$  satisfies condition  $(C_{p'})$ , for  $p' < p$ .

Let  $[0, 1] \setminus \hat{K} =: \sqcup_k U_k$ . For every  $i \geq N_p$  and  $k$ , let  $n_i(k)$  be the number of intervals of the form  $U_{iq}$ , among  $q \in \{0, \dots, 10^i - 1\}$ , contained in  $U_k$ .

We remark that if  $n_i(k) = 1$ , then  $\text{Leb}(U_k) \geq 10^{-i/p}$ , and if  $n_i(k) \geq 2$ , then  $\text{Leb}(U_k) \geq n_i(k)10^{-i}/2$ .

Let us suppose for the sake of contradiction that:

$$\sum_k \text{Leb}(U_k)^{p'} \leq M < \infty$$

This implies that for every  $i$ :

$$(4) \quad \text{Card}\{k : n_i(k) = 1\} \leq M10^{ip'/p}$$

Let  $I$  be an interval of density for  $\hat{K}$ , this means that  $\text{Leb}(\hat{K} \cap I) > 3\text{Leb}(I)/4$ . On the other hand, for  $i$  large, it holds  $\sum_{k: U_k \subset I} n_i(k)10^{-i} \geq 3\text{Leb}(I)/4$ , and so by (4):

$$(5) \quad \sum_{k: U_k \subset I, n_i(k) \geq 2} n_i(k)10^{-i} \geq 3\text{Leb}(I)/4 - M10^{i(\frac{p'}{p}-1)}$$

As  $p' < p$ , for  $i$  large the above sum is greater than  $\text{Leb}(I)/2$ . Hence

$$(6) \quad \sum_{k: U_k \subset I, n_i(k) \geq 2} \text{Leb}(U_k) \geq \sum_{k: U_k \subset I, n_i(k) \geq 2} n_i(k) 10^{-i}/2 \geq \text{Leb}(I)/4$$

A contradiction.

### 4.3 Natural example for which Proposition 0.6 is optimal

We continue with  $K$  as above, made by numbers in  $[0, 1]$  whose decimal expression contains only even numbers. We recall that the Hausdorff dimension of  $K$  is  $\ln(5)/\ln(10)$ .

For every  $i$ , let  $U_i := \{u_{ik} \in [0, 1] : 0 \leq k \leq 10^i - 1\}$  be randomly chosen. With  $j = [i/p]$ , we define:

$$\tilde{K} := [0, 1] \setminus \bigcup_i \bigcup_{0 \leq k \leq 10^i - 1} u_{ik} + [0, 10^{-j}]$$

For every  $v \in U_i$  we write its decimal expression  $v = 0, v_1 \dots v_n \dots$ . Let us study the cardinality of:

$$\{0, v_1 \dots v_n \dots v_j - 0, k_1 \dots k_n \dots k_j : k \in \{0, 2, \dots, 8\}^j, v \in U_i\},$$

This cardinality is greater than  $5^j$  times the cardinality of the following set:

$$E_i := \{(\pi(v_n))_{n \leq j} \in \{0, 1\}^j : v \in U_i\},$$

with  $\pi: \{0, \dots, 9\} \rightarrow \{0, 1\}$  equal to 0 on even numbers and 1 on odd numbers.

The probability for  $E_i$  to do not be equal to  $\{0, 1\}^j$  is:

$$2^i (1 - 1/2^{i/p})^{10^i}$$

Note that if  $p > 1 - d = \ln 2 / \ln 10$ , this probability is equivalent to  $\exp(i \ln 2 - \frac{10^i}{2^{i/p}}) \rightarrow 0$  for  $i$  large.

In other words, if  $(\mathcal{C}_p)$  is not satisfied, then the cardinality of the different intervals  $\{v - k + [0, 10^{-j}] : k \in K, v \in U_i\}$  is equivalent to  $10^j$ . As they are all of length  $10^{-j}$ , they union tends to cover all  $[0, 1]$  when  $i$  is large.

In the case  $p = 1 - d$ , if we fix an element of  $\{0, 1\}^j$ , the probability that it does not belong to  $E_i$  is  $(1 - 1/2^{i/p})^{10^i} = (1 - 1/10^i)^{10^i}$ , which tends to  $1/e$ , and so, in any subinterval  $[a, b]$  of  $[0, 1]$ , the expected number of intervals  $\{v - k + [0, 10^{-j}] : k \in K, v \in U_i\}$  contained in  $[a, b]$  is of the order of  $(1 - 1/e)(b - a)10^j$ , and so, which full probability, the complement union of the arithmetic differences  $v - k + [0, 10^{-j}] : k \in K, v \in U_i, i \geq 1$  has no Lebesgue density points, and so has zero Lebesgue measure.

### 4.4 Example for which Proposition 0.6 is stably close to be optimal

We are going to prove the following result:

**Proposition 4.4.** *For every  $d \in (0, 1)$ , for every  $p > 1 - d$ , there exists a regular Cantor set  $K$  of dimension  $d$ , a Cantor set of positive measure  $\tilde{K}$  which satisfies Condition  $(\mathcal{C}_p)$  and such that for every  $\lambda > 0$ , and for every  $t$ , the set  $\lambda K + t$  is not included in  $\tilde{K}$ .*

**Remark 4.5.** This example is stable by small  $C^2$ -perturbations of  $K$ .

The proof of this Proposition uses the following Theorem.

**Theorem 4.6** ([Mor96], Prop2.1, Theorem II-1.2). *Given  $1 > d, d' > 0$  such that  $d + d' > 1$ , there exist regular Cantor sets  $K, K'$  with Hausdorff dimension respectively  $d, d'$ , that intersect stably. Besides, for any  $\lambda > 0$  and any real number  $t$ ,  $\lambda K + t$  is contained in a gap of  $K'$  or  $K'$  is contained in a gap of  $\lambda K + t$ , or  $(\lambda K + t) \cap K' \neq \emptyset$ . Moreover, this property is  $C^2$ -open in  $(K, K')$ .*

*Proof of Proposition 4.4.* Let  $0 < d, p < 1$  such that  $d + p > 1$ . Let  $K$  and  $K'$  be given by Theorem 4.6 with  $d$  and  $d' < p$ . We can suppose that the convex hulls of  $K$  and  $\tilde{K}$  are  $[0, 1]$ . We are going to construct a compact set  $\hat{K}$  of positive measure included in the following set:

$$X := [0, 1] \setminus \bigcup_{r \in \mathbb{Q} \cap [0, 1]} (K' + r)$$

The set  $X$  has full measure but is not compact. Nevertheless  $X$  cannot contain  $\lambda \cdot K + t$  for every  $\lambda > 0$  and  $t \in \mathbb{R}$ . For the sake of contradiction, suppose this holds. Then  $0 < t \leq 1 - \lambda$ . Let  $r \in \mathbb{Q} \cap (t, t + \lambda)$ . Then, since  $\lambda < 1$ , none of the Cantor sets  $\lambda \cdot K + t, K' + r$  can be contained in a gap of the other, and so, by the above result, they should have non-empty intersection, which contradicts the hypothesis that  $\lambda \cdot K + t$  is contained in  $X$ .

It remains to construct a Cantor set which satisfies condition  $(\mathcal{C}_p)$  and is included in  $X$ .

Let  $M \geq 1$  be large, and for every  $r = p/q \in \mathbb{Q}$  put  $\eta_r := q^{-M}$ . We can now define:

$$\hat{K} := [0, 1] \setminus \bigcup_{r \in \mathbb{Q} \cap [0, 1]} V(K' + r, \eta_r),$$

where  $V(K' + r, \eta_r)$  is the neighborhood of  $K' + r$  made by a covering of it by  $C_{K'} \eta_r^{-d'}$  open intervals of length  $\eta_r$ , with  $C_{K'}$  the  $d'$ -box fuzzy measure. Hence the set  $\hat{K}$  satisfies condition  $(\mathcal{C}_p)$  if:

$$\sum_q q \times C_{K'} \eta_r^{-d'} \eta_r^p < \infty.$$

As  $\eta_r^p = q^{-Mp}$ , the above sum converges if  $1 + Md' - Mp < 0$ , which can be obtained for  $M$  large since  $d' < p$ .

The set  $\hat{K}$  is a compact set of positive Lebesgue measure which does not contains any interval. Let  $\tilde{\hat{K}}$  be the set of condensation point of  $\hat{K}$ , that is  $\hat{K}$  without its points which have a neighbourhood whose intersection with  $\hat{K}$  is countable (which form a countable set). Then  $\tilde{\hat{K}}$  is a Cantor set, satisfies also  $(\mathcal{C}_p)$  (see remark 0.4) and does not contains  $\lambda K + t$  for any  $\lambda > 0$  and  $t$ .  $\square$

## 5 Explanation of the toy model for parameter selection

We recall that if a quadratic map  $f(x) = x^2 + a$  has an absolutely continuous invariant measure (acim) with Lyapunov exponent  $\lambda > 0$ , then for  $C > 0$  small enough,  $\lambda' \in (0, \lambda)$ , the set  $\tilde{K}$  of points  $x$  such that:

$$(\star) \quad \|D_x f^n\| \geq C \lambda'^n, \quad \forall n \geq 0.$$

is a Cantor set of positive Lebesgue measure. The Collet-Eckmann property is that the critical value belongs to such a  $\tilde{K}$  (for a certain  $C, \lambda'$ ). At the neighborhood of such a map, there are good distortion bounds for the transfer phase-parameter spaces, using for instance the puzzle-parapuzzle dictionary. This implies, roughly speaking that the set of parameters  $a$  which satisfy the Collet-Eckmann condition contains a subset which has the same geometry as  $\tilde{K}$ . The set of Collet-Eckmann parameters is in particular of positive measure [BC89, Yoc97]. The support of the acim is also called a non-uniformly hyperbolic attractor.

In higher dimension, the unique paradigmatic (generic) family of maps which has a non-uniformly hyperbolic attractor for a positive set of parameters is the Hénon one's with determinant  $b$  small and parameter  $a \in \mathbb{R}$  [BC89, BY93, Ber11]:

$$f: (x, y) \mapsto (x^2 + a + y, -bx)$$

For such parameters  $a$ , there exists an *SRB probability*  $\mu$  for  $f$ :  $\mu$  is invariant, has a positive Lyapunov exponent, and the conditional measure of  $\mu$  with respect to the unstable manifolds is absolutely continuous.

Let us describe the geometry of these attractors. For such parameters  $a$ , there exists a uniformly hyperbolic horseshoe  $H$ , that can be fairly well approximated by the product of Cantor sets  $K^u \times K^s$  where  $K^u$  has Hausdorff dimension close to 1,  $K^s$  has Hausdorff dimension close to 0, and the unstable direction is almost horizontal. The non-uniformly hyperbolic attractor is equal to closure of the unstable set of  $H$ .

Several constructions of the SRB measure [BV06, Ber11] implies that restricted to any local unstable manifold  $W_{loc}^u(z) \cong [-1, 1] \times \{k\}$ ,  $k \in K^s$ , the set of points of  $W_{loc}^u(z_0)$  which satisfies  $(\star)$  is a Cantor set  $\tilde{K}$  of positive measure for any  $C > 0$  small enough,  $\lambda' \in (0, \lambda)$ . From the fact that the Lyapunov exponent is large with respect to the determinant, every point  $x$  of  $\tilde{K}$  has a long local stable manifold  $W_{loc}^s(x)$ . Moreover the family of curves  $(W_{loc}^s(x))_{x \in \tilde{K}}$  depends Lipschitz on  $x$  in the meaning that there exists a Lipschitz homeomorphism which sends  $\sqcup_{x \in \tilde{K}} W_{loc}^s(x)$  onto  $\tilde{K} \times (-1, 1)$ . Moreover there exists  $C' \in (0, C)$  such that  $z \in \sqcup_{x \in \tilde{K}} W_{loc}^s(x)$  satisfies  $(\star)$  for  $C'$  and  $\lambda$ .

Similarly to the one dimensional case, the idea of the proof [Ber11] was to make (in particular) every the image of  $W_{loc}^u(H)$  by  $f$  tangent to  $\sqcup_{x \in \tilde{K}} W_{loc}^s(x)$ . In other words, for every  $z \in H$ , there exists  $x \in \tilde{K}$  such that  $f(W_{loc}^u(z))$  is tangent to  $W_{loc}^s(x)$ . In particular, there exists a diffeomorphism which sends  $K^s$  into  $\tilde{K}$ .

When the parameter  $a$  vary, the folding points have their  $x$ -coordinate which have nearly the same strictly monotone dependence on  $a$ .

The parameter selection is handle in a same similar way as in dimension 1, and roughly speaking, it is done by including in particular  $K^s$  into a similar Cantor set to  $\tilde{K}$  for a Lebesgue positive set of parameters, for a positive set of translation parameters.

*It is a very rough description* : Actually, the parameter selection is done during an induction which constructs an increasing sequence of unstable curves sets  $([-1, 1] \times K_n^s)_n$  and a decreasing sequence  $(\tilde{K}_n)_n$  of Lebesgue positive subsets of  $[-1, 1]$ , such that  $K_0^s = K^s$  and  $K_n^s$  is close to  $K^s$




Figure 1: Toy model for parameter selection in the Hénon family.

and of Hausdorff dimension small, whereas  $\tilde{K} = \cap_n \tilde{K}_n$ . The construction of  $\tilde{K}_n$  and  $K^s$  depend of the  $n$  first iterates  $f^n(W_{loc}^u(H))$ . Therefore, the geometries of  $\tilde{K}_n$  and  $K_n^s$  are constant only on smaller and smaller parameter intervals. These parameter intervals converge to single points.

Nevertheless, with respect to the above description of non uniformly hyperbolic maps, in order to find new non uniformly hyperbolic attractor with large Hausdorff dimension, the following question seems to be crucial:

## References

- [BC89] Michael Benedicks and Lennart Carleson, *On the Hénon attractor*, IXth International Congress on Mathematical Physics (Swansea, 1988), Hilger, Bristol, 1989, pp. 498–500. MR 1033826 (91m:58097)
- [Ber11] P. Berger, *Abundance of one dimensional non uniformly hyperbolic attractors for surface endomorphisms*, arxiv (2011).
- [BV06] Michael Benedicks and Marcelo Viana, *Random perturbations and statistical properties of Hénon-like maps*, Ann. Inst. H. Poincaré Anal. Non Linéaire **23** (2006), no. 5, 713–752. MR 2259614 (2007f:37042)
- [BY93] M. Benedicks and L.S. Young, *Sinai-bowen-ruelle measures for certain henon maps*, Inventiones Math. (1993), no. 112, 541–576.
- [Mor96] Carlos Gustavo T. de A. Moreira, *Stable intersections of Cantor sets and homoclinic bifurcations*, Ann. Inst. H. Poincaré Anal. Non Linéaire **13** (1996), no. 6, 741–781. MR 1420497 (97h:58114)
- [PT93] Jacob Palis and Floris Takens, *Hyperbolicity and sensitive chaotic dynamics at homoclinic bifurcations*, Cambridge Studies in Advanced Mathematics, vol. 35, Cambridge University Press, Cambridge, 1993, Fractal dimensions and infinitely many attractors. MR 1237641 (94h:58129)
- [Yoc97] J.-C. Yoccoz, *Jackobson’ theorem*, Manuscript (1997).